



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GIBRALTAR.

*Quarantine notice.*GIBRALTAR, *March 9, 1896.*

The board of health this day decided that all vessels from Alexandria shall be subject to quarantine on arrival, unless they have completed twelve days from date of departure and have health on board.

By order,

JOHN C. KING,
Secretary to the Board of Health.

JAPAN.

*One cholera case in Fukuoka Ken.*YOKOHAMA, *March 4, 1896.*

SIR: I have the honor to report that the official returns of the Japanese Home Department show 1 case of cholera as having occurred in the ken of Fukuoka on February 29.

I am, sir, very respectfully,

STUART ELDRIDGE, M. D.,
Sanitary Inspector, U. S. M. H. S.

STATISTICAL REPORTS.

AUSTRALIA—*New South Wales.*—Month of January, 1896. Estimated population, 408,500. Total deaths, 650, including enteric fever, 17; scarlet fever, 2; diphtheria, 2, and whooping cough, 2.

BERMUDA.—Three weeks ended March 20, 1896. Estimated population, 15,013. Total deaths, 2. No deaths from contagious diseases.

CUBA—*Habana.*—Under date of March 28, 1896, the United States sanitary inspector reports as follows:

There were 160 deaths in this city during the week ended March 26, 1896. Two of those deaths were caused by yellow fever, 2 by enteric fever, 6 by so-called pernicious fever, 1 by diphtheria, 5 by enteritis, 6 by smallpox, 16 by pneumonia, and 10 by la grippe.

The 2 deaths by yellow fever during the week occurred in the military hospital.

JAMAICA—*Kingston.*—Month of February, 1896. Estimated population, 19,264. Total deaths, 123, including 1 from diphtheria.

GREAT BRITAIN—*England and Wales.*—The deaths registered in 33 great towns of England and Wales during the week ended March 14 correspond to an annual rate of 20 a thousand of the aggregate population, which is estimated at 10,860,971. The lowest rate was recorded in Croydon, viz, 12.4, and the highest in Sunderland, viz, 24.1 a thousand.

London.—One thousand seven hundred and fifty-two deaths were registered during the week, including measles, 137; scarlet fever, 21; diphtheria, 45; whooping cough, 110; enteric fever, 10, and diarrhea and dysentery, 10. The deaths from all causes corresponded to an annual rate of 20.6 a thousand. In greater London 2,177 deaths were registered,